

# Distributions of atmospheric molecules on the surface of the Saturnian moon, Iapetus

I-Ling Lin(1) and Wing-Huen Ip(2)

(1) Institute of Space Science, National Central University

(2) Institute of Astronomy and Space Science, National Central University

## Abstract

The close flyby observations of Iapetus by the Cassini spacecraft on December 31, 2004, during the Saturn Orbit Insertion have brought many exciting new results. Of particular interest for the atmospheric and surface properties of this icy Saturnian moon has to do with the discovery of CO<sub>2</sub> ice in the form of mixtures in water ice and other substances by the infrared mapping spectrometer. This finding has drawn attention to the possible presence of polar ice caps and a CO<sub>2</sub> exosphere. In addition, the existence of very tenuous atmosphere composed of other species like O<sub>2</sub> and N<sub>2</sub> cannot be ruled out. In this paper we will report on our results from the numerical simulations of the ballistic transport of volatile species on the surface of Iapetus. This work is performed with a view to compare with the measurements of the ion and neutral mass spectrometer during the next close flyby in September 2007.

## References

Buratti, B. J. et al., 2005: Cassini Visual and Infrared Mapping Spectrometer Observations of Iapetus: Detection of CO<sub>2</sub>, *ApJ*, **622**, 149-152.

Brown, R. H. and Matson, D. L., 1987: Thermal effects of insolation propagation into the regoliths of airless bodies, *ICARUS*, **72**, 84-94.

Hodges, R. R., JR., 1973: Helium and Hydrogen in the Lunar Atmosphere, *JGR*, **78**, 8055-8063.

Lebofsky, L. A., 1975: Stability of frosts in the solar system, *ICARUS*, **25**, 205-217.

Palmer, E.E. and Brown, R. H., 2006: Carbon Dioxide Transport on Iapetus, *LPSC*, **37**, 2215.

Spencer, J. R., et al., 2005: Cassini CIRS Observations of Iapetus' Thermal Emission, *Lunar Planet. Sci.*, **36**, 2305.

Cassini-Huygens Home: <http://saturn.jpl.nasa.gov>




Figure 1. Iapetus Thermal Mapping Image. The highest temperature of subsolar point in dark hemisphere reaches ~130 K.

From: <http://saturn.jpl.nasa.gov/science/moons/index.cfm>